arXiv:cond-mat/9802254v2 [cond-mat.supr-con] 15 Feb 2000

Transition Radiation of Moving Abrikosov Vortices.

O.V. Dolgov
Max- Planck-Institut für Festkörperforschung,
Heisenbergstrasse 1, D-70569 Stuttgart, Germany

N. Schopohl
Eberhard-Karls-Universität Tübingen, Institut für Theoretische Physik,
Auf der Morgenstelle 14,
D-72076 Tübingen, Germany

We predict that Abrikosov vortices moving towards the surface of a superconductor emit electromagnetic radiation into free space. Passage of a single vortex line travelling at constant speed $\mathbf{v}$ from a superconducting half space, $x < 0$, towards the vacuum half space, $x > 0$, is discussed in detail. The frequency distribution of the radiated intensity displays a pronounced maximum at microwave frequencies around $\omega \simeq |v_x|/\lambda$, where λ is the magnetic penetration depth. Coherent motion of a lattice of flux lines leads to constructive interference and increases the strength of the radiated power per vortex line by a large factor. Also the radiation source may be used to measure the velocity of Abrikosov vortices and to determine the magnetic penetration depth.

PACS numbers: 41.20.Bt, 41.60.-m, 74.25.Nf, 74.60.-w

A charged particle, moving inside a homogenous medium at constant speed $\mathbf{v}$ and not exceeding the phase velocity c of light in the medium, cannot radiate electromagnetic waves. However, when the particle passes from one medium into another with different electromagnetic properties there may occur transition radiation[1],[2]. In sharp contrast to Cherenkov radiation there exists no lower bound on the velocity of a charged particle for transition radiation.

In the following we suggest that the phenomenon of transition radiation should also occur for electromagnetic objects like Abrikosov vortices moving towards the surface of a superconductor. In our calculation we assume all vortex lines being orientated parallel to the $\widehat{\mathbf{z}}$-axis and we assume $\widehat{\mathbf{z}} \cdot \mathbf{v} = 0$.

Let us first consider a single Abrikosov vortex at rest[35]. Such a vortex is a topological object inside an ordered medium, namely a line defect in a superconductor. Also it is a quantum object. The gradient of the phase of the pairing wave function drives Cooper pairs (i.e. pairs of correlated electrons near the Fermi surface carrying a small center of mass momentum) in circles around the vortex axis, while the modulus of the pairing potential vanishes at the center of the vortex core (say the node is at position $\mathbf{R}$ in the xy-plane). The flow of mass around the vortex axis displays a *quantized* circulation in units of $\frac{h}{2m}$, and the magnetic flux associated with such a line defect is quantized in units of $\Phi_0 = \frac{hc}{2e}$, the quantum of flux. Due to Meissner screening the magnetic field around the vortex decays exponentially fast on the scale of the London penetration depth λ with increasing distance $|\mathbf{r}-\mathbf{R}| \gg \lambda$ to the vortex core:

$$\mathbf{B}(\mathbf{r}) = \widehat{\mathbf{z}}\frac{\Phi_0}{2\pi\lambda^2}\sqrt{\frac{\lambda}{|\mathbf{r}-\mathbf{R}|}}\exp(-\frac{|\mathbf{r}-\mathbf{R}|}{\lambda}) \tag{1}$$

In sharp contrast to this fast decay of the magnetic field $\mathbf{B}(\mathbf{r})$ is the asymptotic behaviour of the vector potential $\mathbf{A}(\mathbf{r})$ around an Abrikosov vortex, which is connected to the magnetic field via $\mathbf{B}(\mathbf{r}) = \operatorname{rot}\mathbf{A}(\mathbf{r})$:

$$\mathbf{A}(\mathbf{r}) = \tfrac{\Phi_0}{2\pi}\,\widehat{\mathbf{z}} \wedge \tfrac{\mathbf{r}-\mathbf{R}}{|\mathbf{r}-\mathbf{R}|^2} \;\text{ for }\; |\mathbf{r}-\mathbf{R}| \gg \lambda \tag{2}$$

Note that $\mathbf{A}(\mathbf{r})$ is independent on λ far outside the London core (apart from exponentially small correction terms).

This slow decay of the vector potential around a single Abrikosov vortex is reminiscent to the behaviour of the vector potential in the exterior region of a long solenoid, like in an *Aharonov-Bohm* experiment. Consider a solenoid consisting out of a piece of infinitesimally thin wire wound helically around the axis $\widehat{\mathbf{z}}$ of an infinitely long cylinder with radius r_σ. Through the wire of the solenoid flows a suitable electrical current I_σ to generate a constant magnetic field $\mathbf{B}^{(\sigma)} = B^{(\sigma)}\widehat{\mathbf{z}}$ *inside*, but *outside* the solenoid the magnetic field is zero. The vector potential $\mathbf{A}^{(\sigma)}$ generated by the solenoid in its exterior region is the gradient of a 'phase':

$$\mathbf{A}^{(\sigma)}(\mathbf{r}) = \begin{cases} \frac{B^{(\sigma)}}{2}\widehat{\mathbf{z}} \wedge (\mathbf{r}-\mathbf{R}) & \text{for} \quad |\mathbf{r}-\mathbf{R}| \le r_\sigma \\ \frac{\Phi_\sigma}{2\pi}\,\nabla\vartheta\,(\mathbf{r}-\mathbf{R}) & \text{for} \quad |\mathbf{r}-\mathbf{R}| > r_\sigma \end{cases} \tag{3}$$

where $e^{i\vartheta(\mathbf{r})} = \frac{x+iy}{\sqrt{x^2+y^2}}$, $\nabla\vartheta(\mathbf{r}) = \widehat{\mathbf{z}} \wedge \frac{\mathbf{r}}{|\mathbf{r}|^2}$ and $\mathbf{r} = (x, y)$. Consider a closed path Γ in the exterior region of the solenoid with winding number 1 around the $\widehat{\mathbf{z}}-$axis. Integration of $\mathbf{A}^{(\sigma)}(\mathbf{r})$ along Γ leads to:

$$\oint_\Gamma d\mathbf{r}\cdot\mathbf{A}^{(\sigma)}(\mathbf{r}) = \Phi_\sigma = \int_\Omega d^2r\;\widehat{\mathbf{z}}\cdot\operatorname{rot}\mathbf{A}^{(\sigma)}(\mathbf{r}) = \pi r_\sigma^2\,B^{(\sigma)} \tag{4}$$

(apply Stokes theorem to the area Ω enclosed by Γ) . So, the constant Φ_σ is equal to the flux passing through the solenoid. Adjusting the control current I_σ such that Φ_σ becomes equal to one quantum of flux, $|\Phi_\sigma| = \Phi_0$, and taking the limit $r_\sigma \to 0^+$, the thin solenoid produces (in its rest frame) a singular electromagnetic field at its center:

$$\begin{aligned} \mathbf{B}^{(\sigma)}(\mathbf{r}) &= \Phi_\sigma \widehat{\mathbf{z}}\, \delta^{(2)}(\mathbf{r}-\mathbf{R}) \\ \mathbf{E}^{(\sigma)}(\mathbf{r}) &= \mathbf{0} \end{aligned} \tag{5}$$

Consider now a thin solenoid travelling relative to the lab frame at a constant speed $\mathbf{v}$ along the trajectory $\mathbf{R} = \mathbf{v}t$. Such a solenoid is surrounded by a *time dependent* Aharonov-Bohm type vector potential giving rise not only to a singular magnetic field $\mathbf{B}^{(\sigma)}(\mathbf{r}, t)$ but also to a singular electric field $\mathbf{E}^{(\sigma)}(\mathbf{r}, t)$:

$$\begin{aligned} \mathbf{B}^{(\sigma)}(\mathbf{r},t) &= \Phi_\sigma \widehat{\mathbf{z}}\, \delta^{(2)}(\mathbf{r}-\mathbf{v}t) \\ \mathbf{E}^{(\sigma)}(\mathbf{r},t) &= -\frac{\mathbf{v}}{c} \wedge \mathbf{B}^{(\sigma)}(\mathbf{r},t) \end{aligned} \tag{6}$$

It is consistent (and convenient) to associate with the moving thin solenoid a *conserved* charge-current density via:

$$\begin{aligned} 4\pi\rho^{(\sigma)}(\mathbf{r},t) &= \operatorname{div} \mathbf{E}^{(\sigma)}(\mathbf{r},t) \\ \frac{4\pi}{c}\mathbf{J}^{(\sigma)}(\mathbf{r},t) &= \operatorname{rot} \mathbf{B}^{(\sigma)}(\mathbf{r},t) - \frac{1}{c}\partial_t \mathbf{E}^{(\sigma)}(\mathbf{r},t) \end{aligned} \tag{7}$$

Since in the geometry under consideration translational invariance is broken at $x = 0$ it is natural to require that no current must flow from the region $x < 0$ (superconductor) into the region $x > 0$ (vacuum):

$$\lim_{x\to 0} \widehat{\mathbf{n}} \cdot \mathbf{J}^{(\sigma)}(\mathbf{r},t) = 0 \tag{8}$$

($\widehat{\mathbf{n}}$ denotes the normal unit vector of the interface). By a simple trick we are able to fulfill this boundary condition (which actually avoids solving an integral equation problem of the Wiener-Hopf type). Let us superpose to the field around a single flux line situated at its (instantaneous) position $\mathbf{R} = \mathbf{v}t$ another singular field generated by a mirror vortex (with *opposite* magnetic moment) situated at the *mirror reflected* position $\widetilde{\mathbf{R}} = \widetilde{\mathbf{v}}t$, where $\widetilde{\mathbf{v}} = (-v_x, v_y)$. The resulting fields (inside the region $x < 0$) are then:

$$\begin{aligned} \mathbf{B}^{(\sigma)}(\mathbf{r},t) &= \Phi_\sigma \widehat{\mathbf{z}} \left[\delta^{(2)}(\mathbf{r}-\mathbf{v}t) - \delta^{(2)}(\mathbf{r}-\widetilde{\mathbf{v}}t)\right] \\ \mathbf{E}^{(\sigma)}(\mathbf{r},t) &= \Phi_\sigma \widehat{\mathbf{z}} \wedge \left[\frac{\mathbf{v}}{c}\delta^{(2)}(\mathbf{r}-\mathbf{v}t) - \frac{\widetilde{\mathbf{v}}}{c}\delta^{(2)}(\mathbf{r}-\widetilde{\mathbf{v}}t)\right] \end{aligned} \tag{9}$$

Inserting these expressions for $\mathbf{B}^{(\sigma)}$ and $\mathbf{E}^{(\sigma)}$ into Eq.(7) there results a *conserved* charge-current distribution which generates the asymptotics of the Aharonov-Bohm type vector potential around a *moving* Abrikosov vortex line far outside the London core region. By construction this charge-current distribution also fulfills the boundary value condition Eq.(8).

Let us next discuss the effect of Meissner screening. Since a superconductor is an *ordered* medium it will react to a specified external influence by a specific response that takes into account the formation of induced currents and of induced polarization charges inside the sample. This means a *parametrically* driven flux line moving at speed $\mathbf{v}$ inside a medium with Meissner screening (imagine an external agent who pulls the solenoid along) is surrounded by *total* current and charge densities of the form

$$\begin{aligned} \mathbf{J}^{(tot)}(\mathbf{r},\omega) &= \mathbf{J}^{(ind)}(\mathbf{r},\omega) + \mathbf{J}^{(\sigma)}(\mathbf{r},\omega) \\ \rho^{(tot)}(\mathbf{r},\omega) &= \rho^{(ind)}(\mathbf{r},\omega) + \rho^{(\sigma)}(\mathbf{r},\omega) \end{aligned} \tag{10}$$

Here $\mathbf{J}^{(ind)}$ and $\rho^{(ind)}$ represent the induced response of the ordered medium to the applied drive. Explicit expressions for the drive terms $\mathbf{J}^{(\sigma)}(\mathbf{r},\omega)$ and $\rho^{(\sigma)}(\mathbf{r},\omega)$ follow inserting Eq.(9) into Eq.(7) and calculating the Fourier transforms with respect to time. Since the induced response is retarded in time it is convenient to switch to the frequency representation. Suitable expressions for $\rho^{(ind)}(\mathbf{r},\omega)$ and $\mathbf{J}^{(ind)}(\mathbf{r},\omega)$ follow from standard linear response theory. Since any *local* perturbation inside the bulk of a superconductor heals typically on the length scale of the coherence length ξ of the Cooper pairs ($\xi \ll \lambda$ in extreme type-II materials), and since the phenomenon we study is related to low frequencies in the microwave regime, a suitable (local) expression for the induced charge-current distribution is:

$$\begin{aligned} \mathbf{J}^{(ind)}(\mathbf{r},\omega) &= -\frac{c}{4\pi}\,\frac{c}{i\omega}\frac{1}{\lambda_\omega^2}\,\mathbf{E}^{(tot)}(\mathbf{r},\omega) \\ \rho^{(ind)}(\mathbf{r},\omega) &= \frac{1}{i\omega}\operatorname{div}\mathbf{J}^{(ind)}(\mathbf{r},\omega) \end{aligned} \tag{11}$$

In a *'two fluid'* type description of the electromagnetic response of a superconductor it is found that the length λ_ω is a frequency dependent (but local) penetration depth, that interpolates between the skin penetration depth, which is characteristic for the normal metallic state at frequencies above 2Δ, to the magnetic penetration depth (MPD) at zero frequency:

$$\frac{1}{\lambda_\omega^2} = \frac{\omega_{pl}^2}{c^2}\frac{\omega}{\omega + i\gamma}\Theta(|\omega| - 2\Delta) + \frac{1}{\lambda^2}\Theta(2\Delta - |\omega|) \tag{12}$$

Here $\omega_{pl} = \sqrt{\frac{4\pi n\, e^2}{m^*}}$ is the bulk plasma frequency, γ describes quasiparticle damping, $\lambda = \lambda(\gamma)$ denotes MPD of a superconductor[6]. In the clean limit (and at zero temperature) $\lim_{\gamma\to 0}\lambda(\gamma) = \frac{c}{\omega_{pl}} = \lambda_L$, the standard definition of the London penetration depth. Of course, the nonlocality of the electromagnetic response kernel becomes eventually important very near to the surface (e.g. anomalous skin effect).

In the presence of Meissner screening the (total) electromagnetic field around a travelling thin solenoid consists out of *two* parts. The *first* part is entirely due to the Meissner screening currents which flow in the exterior region of the solenoid. This part actually represents (within the accuracy of local London electrodynamics) the electromagnetic fields around a moving Abrikosov vortex, provided the flux inside the solenoid is fixed with the opposite sign: $\Phi_\sigma = -\Phi_0$. The *second* part originates from the infinitesimal small *interior* region of the thin solenoid, where there exists (in the rest frame of the solenoid) an artificially fixed singular magnetic field: $B^{(\sigma)} = \frac{\Phi_\sigma}{\pi r_\sigma^2}$. This feature of the thin solenoid model is in sharp contrast to what really happens at the center of an Abrikosov vortex, where the pairing wave function displays a node and where the magnetic field assumes a finite and (almost) constant value inside the inner core of the vortex , a region of size $\xi \ll \lambda$. As a result, the self fields $\mathbf{E}^{(\sigma)}(\mathbf{r},\omega)$ and $\mathbf{B}^{(\sigma)}(\mathbf{r},\omega)$, which originate from the *interior* region of the solenoid, should be subtracted from the total fields $\mathbf{E}^{(tot)}(\mathbf{r},\omega)$ and $\mathbf{B}^{(tot)}(\mathbf{r},\omega)$ in order to determine the physical fields $\mathbf{E}(\mathbf{r},\omega)$ and $\mathbf{B}(\mathbf{r},\omega)$ around an Abrikosov vortex line:

$$\begin{aligned} \mathbf{B}^{(tot)}(\mathbf{r},\omega) &= \mathbf{B}(\mathbf{r},\omega) + \mathbf{B}^{(\sigma)}(\mathbf{r},\omega) \qquad (13) \\ \mathbf{E}^{(tot)}(\mathbf{r},\omega) &= \mathbf{E}(\mathbf{r},\omega) + \mathbf{E}^{(\sigma)}(\mathbf{r},\omega) \end{aligned}$$

The fields $\mathbf{E}^{(tot)}(\mathbf{r},\omega)$ and $\mathbf{B}^{(tot)}(\mathbf{r},\omega)$ are solutions to the Maxwell equations in the presence of the total charge-current distribution specified in Eq.(10):

$$\begin{aligned} &\frac{4\pi}{c}\left[\mathbf{J}^{(\sigma)}(\mathbf{r},\omega) + \mathbf{J}^{(ind)}(\mathbf{r},\omega)\right] \qquad (14) \\ &= \operatorname{rot}\mathbf{B}^{(tot)}(\mathbf{r},\omega) + \frac{i\omega}{c}\mathbf{E}^{(tot)}(\mathbf{r},\omega) \end{aligned}$$

Upon insertion of Eq.(11) this transforms into:

$$\begin{aligned} &\frac{4\pi}{c}\mathbf{J}^{(\sigma)}(\mathbf{r},\omega) - \frac{c}{i\omega}\frac{1}{\lambda_\omega^2}\left[\mathbf{E}(\mathbf{r},\omega) + \mathbf{E}^{(\sigma)}(\mathbf{r},\omega)\right] \qquad (15) \\ &= \operatorname{rot}\left[\mathbf{B}(\mathbf{r},\omega) + \mathbf{B}^{(\sigma)}(\mathbf{r},\omega)\right] + \frac{i\omega}{c}\left[\mathbf{E}(\mathbf{r},\omega) + \mathbf{E}^{(\sigma)}(\mathbf{r},\omega)\right] \end{aligned}$$

Next we insert $\mathbf{J}^{(\sigma)}$ from the definition Eq.(7) and obtain:

$$-\frac{c}{i\omega}\frac{1}{\lambda_\omega^2}\left[\mathbf{E}(\mathbf{r},\omega) + \mathbf{E}^{(\sigma)}(\mathbf{r},\omega)\right] = \operatorname{rot}\mathbf{B}(\mathbf{r},\omega) + \frac{i\omega}{c}\mathbf{E}(\mathbf{r},\omega) \qquad (16)$$

Using the Faraday law

$$\operatorname{rot}\mathbf{E}(\mathbf{r},\omega) - \frac{i\omega}{c}\mathbf{B}(\mathbf{r},\omega) = \mathbf{0}$$

it is not difficult to derive *dynamical* London equations for the physical electromagnetic fields around a moving vortex line:

$$\begin{aligned} \operatorname{rot}\operatorname{rot}\mathbf{E}(\mathbf{r},\omega) + \left(\frac{1}{\lambda_\omega^2} - \frac{\omega^2}{c^2}\right)\mathbf{E}(\mathbf{r},\omega) &= \frac{-1}{\lambda_\omega^2}\,\mathbf{E}^{(\sigma)}(\mathbf{r},\omega) \qquad (17) \\ \left[-\mathbf{\Delta} + \left(\frac{1}{\lambda_\omega^2} - \frac{\omega^2}{c^2}\right)\right]\mathbf{B}(\mathbf{r},\omega) &= \frac{-1}{\lambda_\omega^2}\,\mathbf{B}^{(\sigma)}(\mathbf{r},\omega) \end{aligned}$$

For $x > 0$ we have $\frac{1}{\lambda_\omega^2} \equiv 0$, of course, since there is nothing in vacuum. In the *static* limit, $|\mathbf{v}| \to 0$ and $\omega \to 0$, the second equation reduces to the standard London equation determining the magnetic field around an Abrikosov vortex line at rest[4],[3],[5]. Note that for simplicity we assume here that there exists no external static magnetic field in vacuum, i.e. $\lim_{\omega\to 0}\mathbf{B}(\mathbf{r},\omega) = \mathbf{0}$ for $x > 0$.

Next we discuss scattering solutions to our dynamical London equations, which describe the transition radiation effect for moving Abrikosov vortices. Before we expand details of the new radiation source let us comment on an interesting difference to standard transition radiation[1],[2]. In the case of standard transition radiation the moving particle, say an electron or a proton, actually exists on either side of the interface passing from one medium into another with different electromagnetic properties. In the case of a moving vortex the 'particle' (i.e.the travelling singular line representing the center of the vortex), exists only inside the superconductor, but ceases to exist on the other side of the interface (in vacuum).

A Fourier transformation of the dynamical London equations with respect to the $y-$coordinate simplifies the problem and leads to coupled ordinary differential equations. It is sufficient to consider the tangential component $E_y(x) = E_y(x, k_y, \omega)$ of the electric field, since the normal component $E_x(x) = E_x(x, k_y, \omega)$ follows from knowledge of $\partial_x E_y(x)$. From both components of the electric field the magnetic field $B_z(x) = B_z(x, k_y, \omega)$ may be calculated using Faraday's law. Guided by the structure of our problem we make now the following ansatz for $E_y(x)$ in terms of three unknown coefficients a_S , b_S and a_V , and in terms of three (complex) wave numbers q_λ , q_∞ and K_ω:

$$E_y(x) = \begin{cases} a_S \exp\left(q_\lambda x\right) + b_S\left(e^{iK_\omega x} + e^{-iK_\omega x}\right) & \text{for } \; x < 0 \\ a_V \exp\left(-q_\infty x\right) & \text{for } \; x > 0 \end{cases} \qquad (18)$$

$$\begin{aligned} q_\infty &= \sqrt{k_y^2 - \frac{\omega^2}{c^2}}\;,\; q_\lambda = \sqrt{k_y^2 + \frac{1}{\lambda_\omega^2} - \frac{\omega^2}{c^2}} \\ K_\omega &= \frac{\omega - k_y v_y}{v_x} \qquad (19) \end{aligned}$$

At infinity ($|x| \to \infty$) we pose the usual radiation boundary conditions. So, in order to avoid exponential growth for $x \to \pm\infty$, the branch cut of the square root is defined such that $\operatorname{Re} q_\lambda > 0$ and $\operatorname{Re} q_\infty > 0$. The coefficient b_S follows from the requirement that $E_y(x)$ should be a solution to the differential equation in the half space $x < 0$. The coefficients a_S and a_V follow from the standard boundary value conditions for the tangential and normal components of the electric field at $x = 0$. The results are:

$$b_S = \frac{\operatorname{sign}(v_x)}{c}\,\frac{\Phi_0}{\lambda_\omega^2}\,\frac{\left(\frac{v_x}{c}q_\lambda\right)^2 - \frac{v_y}{c}\,k_y\,K_\omega}{\frac{1}{\lambda_\omega^2} - \frac{\omega^2}{c^2}}\,\frac{1}{K_\omega^2 + \left|\frac{v_x}{c}\right|^2 q_\lambda^2}$$

$$
\begin{aligned}
a_S &= \frac{2b_S\,\frac{\omega^2}{c^2}\,q_\lambda}{\left(-\frac{\omega^2}{c^2}\right)q_\lambda + \left(\frac{1}{\lambda_\omega^2} - \frac{\omega^2}{c^2}\right)q_\infty} \\
a_V &= \frac{2b_S\,\left(\frac{1}{\lambda_\omega^2} - \frac{\omega^2}{c^2}\right)\,q_\infty}{\left(-\frac{\omega^2}{c^2}\right)q_\lambda + \left(\frac{1}{\lambda_\omega^2} - \frac{\omega^2}{c^2}\right)q_\infty}
\end{aligned} \tag{20}
$$

The coefficient b_S displays a resonance around $\omega = \omega_{pl}$, while the coefficients a_S and a_V display also a resonance around $\omega = \frac{\omega_{pl}}{\sqrt{2}}$, the surface plasma frequency. The solution for $E_y(x)$ in vacuum ($x > 0$) describes an outward travelling wave front, representing electromagnetic radiation. The radiation part is entirely determined by oscillating terms, which exist only in the range of wave numbers $|k_y| < \left|\frac{\omega}{c}\right|$, so that $q_\infty = -i\,\mathrm{sign}(\omega)\sqrt{\frac{\omega^2}{c^2} - k_y^2}$ is purely imaginary. For $|k_y| > \left|\frac{\omega}{c}\right|$ the modulus of the electromagnetic field in vacuum is exponentially small for $x >> \frac{c}{|\omega|}$, and may be safely neglected.

Consider now a planar detector screen at a distance $x_T \gg \frac{c}{|\omega|}$ in the far zone in vacuum, with the plane of the screen orientated parallel to the superconductor-vacuum interface (at $x = 0$). The rate at which the emitted radiation is incident onto a target area element $\widehat{\mathbf{n}}\ da = -\widehat{\mathbf{x}}\ da$ at position $\mathbf{r}_T = (x_T, y_T)$ on the screen may be calculated from the associated Poynting flux $\frac{c}{4\pi}\ [\mathbf{E}(\mathbf{r}_T, t) \wedge \mathbf{B}(\mathbf{r}_T, t)]\cdot(-\widehat{\mathbf{x}})$. Let us denote the amount of energy $d\varepsilon(\mathbf{r}_T)$ so radiated (per length unit L of the vortex line) onto an infinitesimal target area element $da_T = L \cdot dy_T$ by $\frac{d\varepsilon(\mathbf{r}_T)}{da_T}$. Then the total energy ε_{rad} radiated into free space (per unit length L of the vortex line) is obtained integrating $\frac{d\varepsilon(\mathbf{r}_T)}{da_T}$ over time and over the the entire screen area (wide detector):

$$
\frac{\varepsilon_{rad}}{L} = \frac{c}{4\pi}\int_{-\infty}^{\infty}\frac{d\omega}{2\pi}\int_{-\left|\frac{\omega}{c}\right|}^{\left|\frac{\omega}{c}\right|}\frac{dk_y}{2\pi}\,E_y(x_T, k_y, \omega)B_z(x_T, -k_y, -\omega)
$$

Let us introduce scaled variables: $\overline{\omega} = \frac{\omega}{\omega_{pl}}$, $\overline{\lambda} = \frac{\lambda(\gamma)}{\lambda_L}$, $\overline{k}_y = \lambda_L k_y$, $\overline{q}_\infty = \lambda_L q_\infty$, $\overline{q}_\lambda = \lambda_L q_\lambda$ and $\overline{v}_x = \frac{v_x}{c}$, $\overline{v}_y = \frac{v_y}{c}$, so that energy is measured in units of $\hbar\omega_{pl}$, length is measured in units of $\lambda_L = \frac{c}{\omega_{pl}}$ and velocity is measured in units of c. We may then rewrite the vacuum amplitude $a_V = a(k_y, \omega) = \frac{\Phi_0}{c}\,A_V\left(\overline{k}_y, \overline{\omega}\right)$ in dimensionless form:

$$
A_V = \frac{2}{\overline{\lambda}_\omega^2}\,\frac{\mathrm{sign}(v_x)}{\left(\overline{\omega} - \overline{v}_y\overline{k}_y\right)^2 + \overline{v}_x^2\overline{q}_\lambda^2}\,\frac{\overline{q}_\infty\left(\overline{v}_x^2\overline{q}_\lambda^2 + \overline{v}_y^2\overline{k}_y^2 - \overline{v}_y\overline{k}_y\overline{\omega}\right)}{\left(\frac{1}{\overline{\lambda}_\omega^2} - \overline{\omega}^2\right)\overline{q}_\infty - \overline{\omega}^2\overline{q}_\lambda} \tag{21}
$$

It follows

$$
\frac{\varepsilon_{rad}}{L} = \left(\frac{\Phi_0}{2\pi\lambda_L}\right)^2\frac{1}{2}\int_{-\frac{\pi}{2}}^{\frac{\pi}{2}}\frac{d\theta}{\pi}\int_0^\infty d\overline{\omega}\ \overline{\omega}\ \ \left|A_V\left(\overline{\omega}\sin\theta, \overline{\omega}\right)\right|^2 \tag{22}
$$

The function $I(\theta, \overline{\omega}) = \overline{\omega} \cdot \left|A_V\left(\overline{\omega}\sin\theta, \overline{\omega}\right)\right|^2$ is the angle-frequency distribution of transition radiation emitted into free space by a single moving Abrikosov vortex. Apparently, the dependence of $I(\theta, \overline{\omega})$ on the angle θ for a non relativistic speed is such that it is practically constant for all directions, with the exception of grazing angles θ around $|\theta| \lessapprox \frac{\pi}{2}$. In Fig.(1) we plot the radiated intensity $I(\overline{\omega}) = \int_{-\frac{\pi}{2}}^{\frac{\pi}{2}}\frac{d\theta}{\pi}I(\theta, \overline{\omega})$ vs. scaled frequency $\overline{\omega}$ for various vortex speeds. Values for the gap, the quasiparticle damping γ and the vortex speeds are taken as indicated in the Figure caption. The radiated intensity $I(\overline{\omega})$ displays a sharp peak at a frequency

$$
2\pi\nu_{peak} = \omega_{peak} = \frac{1}{\sqrt{3}}\frac{|v_x|}{\lambda(\gamma)} \tag{23}
$$

This is in agreement with the physical picture, that it takes a time $T \simeq \frac{2\lambda_L}{|v_x|}$ for the London core of a vortex to pass the interfacial region. Our analytical estimatation for ω_{peak} applies provided the frequency of the emitted radiation fulfills $\hbar\omega < 2\Delta$. Indeed, even for a very high vortex speed around $|v_x| \simeq v_F$ on finds $\hbar\omega_{peak} = \frac{1}{\sqrt{3}}\frac{\hbar v_F}{\lambda(\gamma)} \simeq \frac{\xi_{BCS}}{\lambda_L}\cdot\Delta \ll \Delta$, since $\xi_{BCS} \simeq \frac{\hbar v_F}{\Delta} \ll \lambda_L$. Rather high vortex speeds around $v_x \simeq 10^3\frac{m}{\mathrm{sec}}$ have been measured in layered type-II materials in the experiments of Doettinger et al.[7]. For a London penetration depth of order $\lambda(\gamma) \simeq 10^3$Å we expect the position of the peak at $\nu_{peak} = \frac{\omega_{peak}}{2\pi} \simeq \frac{1}{\sqrt{3}2\pi}\frac{10^3\frac{m}{\mathrm{sec}}}{10^3\text{Å}} \simeq 1\,GHz$, a typical microwave frequency. Also the height of the peak intensity increases linearly with the vortex speed v_x, as shown in Fig.(1), in agreement with our analytical estimate $I(\overline{\omega}_{peak}) \simeq \frac{3}{4}\sqrt{3}\,\frac{\overline{v}_x}{\overline{\lambda}}$. Finally, the total amount of energy that gets emitted into free space due to the transition radiation effect follows integrating over all frequencies. Certainly, at very high frequencies, say for $\omega > \omega_{pl}$, our local electrodynamics becomes invalid. But since the integral converges rapidly the error is small if we simply extend the upper integration limit towards infinity. We obtain:

$$
\frac{\varepsilon_{rad}}{L} \simeq \left[\frac{\Phi_0}{2\pi\lambda(\gamma)}\right]^2\left(\frac{v_x}{c}\right)^2 \tag{24}
$$

Assuming $\lambda_L \simeq 100\,nm$ and a vortex speed of order $\frac{v_x}{c} \simeq 10^{-5}$ gives $\varepsilon_{rad} \simeq 10^{-22}\,W \cdot sec$ (per length $L = 1\,cm$). Due to the large impedance mismatch of the superconductor with the vacuum ($Z_0 \simeq 377\,\Omega$) it is not surprising that only a tiny fraction of the nucleation energy[3],[4] of a single Abrikosov vortex gets radiated into free space. So, when the line is destroyed during its passage through the interfacial region, the energy that is not radiated into free space gets converted into elementary excitations (e.g. polaritons) and, eventually, into heat.

A larger effect arises when several vortices, say a bundle of altogether $|\Lambda|$ flux lines, move coherently at constant speed $\mathbf{v}$ towards the surface of a superconductor. According to the superposition principle the (effective) drive term for several flux lines, situated at time $t = 0$ at positions $\mathbf{r}_L = (x_L, y_L)$ inside the superconductor, follows from Eq.(9) upon summation over the instantaneous positions $\mathbf{R}_L = \mathbf{r}_L + \mathbf{v}\ t$ of the individual vortices (and the instantaneous positions of the mirror vortices at $\widetilde{\mathbf{R}}_L = \widetilde{\mathbf{r}}_L + \widetilde{\mathbf{v}}\ t$). The linearity of the dynamical London equations Eq.(17) implies that the electromagnetic field $\mathbf{E}^{(\Lambda)}$ generated by such a bundle of flux lines is:

$$\mathbf{E}^{(\Lambda)}(x, k_y,\omega) = \sum_{\mathbf{r}_L \epsilon \Lambda} e^{-i(k_y y_L + K_\omega x_L)} \mathbf{E}(x, k_y,\omega) \qquad (25)$$

From the theory of diffraction of electromagnetic waves it follows that the associated Poynting flux, and hence the radiated peak intensity emitted by a *periodic* arrangement of coherently moving flux lines (for example, a square vortex lattice Λ, with lattice constant d_Λ, consisting out of $|\Lambda| = N_x \cdot N_y$ flux lines), should be increased by a factor proportional to the number of participating modes compared to the case where only unlocked modes are superposed. The total radiated energy (per unit length L of the vortex lines) is

$$\frac{\varepsilon_{rad}^{(\Lambda)}}{L} = \int_{-\infty}^{\infty} \frac{d\omega}{2\pi} \frac{c}{4\pi} \int_{-\left|\frac{\omega}{c}\right|}^{\left|\frac{\omega}{c}\right|} \frac{dk_y}{2\pi} \frac{\left|\frac{\omega}{c}\right| \, |a_V\,(k_y,\omega)|^2 \, |S_\Lambda(k_y,\omega)|^2}{\sqrt{\frac{\omega^2}{c^2} - k_y^2}}$$

The structure factor of the moving vortex lattice

$$S_\Lambda(k_y, \omega) = \left| \sum_{\mathbf{r}_L \epsilon \Lambda} e^{-i(k_y\, y_L + K_\omega x_L)} \right|^2 \qquad (26)$$

represents, for large N_x and N_y, a series of delta spikes, selecting wave numbers k_y and frequencies ω such that $k_y = \frac{2\pi}{d_\Lambda} n_y$ and $\omega = \frac{2\pi}{d_\Lambda}(n_x v_x + n_y v_y)$, with n_x and n_y integer. For intermediate magnetic fields $B_{c1} \ll B \ll B_{c2}$ the lattice constant d_Λ is much larger than the coherence length ξ, but smaller than the London depth λ_L , so that $\xi \ll d_\Lambda \ll \lambda_L$. We derive from this a simple estimation for the radiation energy emitted into free space by alltogether $|\Lambda| \gg 1$ coherently moving vortex lines (arranged in the geometry of a square lattice as described above):

$$\frac{\varepsilon_{rad}^{(\Lambda)}}{L} \simeq \frac{|\Lambda|}{\pi} \left(\frac{\Phi_0}{d_\Lambda}\right)^2 \frac{|v_x|}{c} \qquad (27)$$

The principal result is that the transition radiation effect for a cluster of altogether $|\Lambda| \gg 1$ *coherently* moving flux lines is a *linear* function of the vortex speed $|v_x|$. The emitted power *per vortex* is huge compared to the radiation emitted by a single vortex by a factor proportional to the number of vortices, since $\frac{1}{d_\Lambda^2} \propto |\Lambda|$. This enhancement is due to *constructive* interference, like in the super radiation effect of laser physics. Assuming $\lambda_L \simeq 100\, nm$ and $d_\Lambda = 0.5\, \lambda_L$ gives a total of $|\Lambda| \simeq 0.25 \times 10^{12}$ vortices per cm^2. This leads (per length $L = 1\, cm$) and a vortex speed $v_x \simeq 10^{-5} c$, $v_y = 0$ to a value of order $\varepsilon_{rad}^{(\Lambda)} \simeq 0.12 \times 10^{-3}\, W\, sec$. This estimation suggests that a sensitive S-I-S Josephson radiometer could be used to detect the predicted spectrum of transition radiation in free space. The sharp peak of the radiated intensity around $\omega \sim \frac{|v_x|}{\lambda}$ allows to measure MPD for a known vortex speed, and vice versa.

Our calculations indicate that transition radiation is a *generic* phenomenon, that should exist in other types of *ordered* condensed matter containing mobile (topological) defects. We expect transition radiation to be emitted into free space by rapidly moving dislocations, by magnetic bubbles and Bloch walls in magnets, and also by travelling polarons.

Acknowledgments: We thank V.L. Ginzburg and A.A. Abrikosov for the interest they have shown into our work and for correspondence. It is a pleasure to thank D. Rainer for an illuminating discussion of the thin solenoid model, and to thank R.P. Huebener for useful information on high speed flux-flow experiments.

---

[1] V.L. Ginzburg and I.M. Frank, Zh. Eksp. Teor. Fiz. **16**, 15 (1946).

[2] V.L. Ginzburg and V.N. Tsytovich, 'Transition Radiation and Transition Scattering' , (Adam Hilger, New York, 1990).

[3] A.A. Abrikosov, "Fundamentals of the Theory of Metals", (North-Holland, Amsterdam, 1988).

[4] P.G. de Gennes,"Superconductivity of Metals and Alloys",(Addison-Wesley, New York, 1989).

[5] V.V. Shmidt and G.S. Mkrtchyan, Usp. Fiz. Nauk **112**, 459 (1974); Sov. Phys.-Usp. **17**, 170 (1974).

[6] Sang Boo Nam, Physical Review **156**, 470 (1967).

[7] S.G. Doettinger, S. Kittelberger, R.P. Huebener and C.C. Tsuei, Phys. Rev. B **56**, 14157 (1997).

FIG. 1. Frequency distribution $I(\overline{\omega})$ of intensity of transition radiation emitted into free space for different vortex speeds; the labels $a), b), c), d)$ correspond to $v_y = 0$ and to $v_x/c = 2 \times 10^{-5}, 1.5 \times 10^{-5}, 10^{-5}, 5 \times 10^{-6}$, respectively. Assumed values for scattering rate $\gamma/\omega_{pl} = 10^{-4}$ and gap $\Delta/\omega_{pl} = 10^{-3}$.


x 10^−5
2.5
2
1.5
1
0.5
0
I(ω)
a)
b)
c)
d)
10^−7
10^−6
10^−5
10^−4
ω/ω_pl